

# Benjamin Sambale

TU Kaiserslautern

Research Member (GRTA)

# Benjamin Sambale

TU Kaiserslautern

Research Member (GRTA)

Academic genealogy:

... → Gauß → Bessel → Scherk → Kummer → Frobenius → Schur  
→ Brauer → Fong → Olsson → Külshammer → me → Dauter?

# Benjamin Sambale

TU Kaiserslautern

Research Member (GRTA)

Academic genealogy:

... → Gauß → Bessel → Scherk → Kummer → Frobenius → Schur  
→ Brauer → Fong → Olsson → Külshammer → me → Dauter?

My preferences:

# Benjamin Sambale

TU Kaiserslautern

Research Member (GRTA)

Academic genealogy:

... → Gauß → Bessel → Scherk → Kummer → Frobenius → Schur  
→ Brauer → Fong → Olsson → Külshammer → me → Dauter?

My preferences:

groups

algebras

# Benjamin Sambale

TU Kaiserslautern

Research Member (GRTA)

Academic genealogy:

... → Gauß → Bessel → Scherk → Kummer → Frobenius → Schur  
→ Brauer → Fong → Olsson → Külshammer → me → Dauter?

My preferences:

groups

algebras

# Benjamin Sambale

TU Kaiserslautern

Research Member (GRTA)

Academic genealogy:

... → Gauß → Bessel → Scherk → Kummer → Frobenius → Schur  
→ Brauer → Fong → Olsson → Külshammer → me → Dauter?

My preferences:

groups

finite

algebras

infinite

# Benjamin Sambale

TU Kaiserslautern

Research Member (GRTA)

Academic genealogy:

... → Gauß → Bessel → Scherk → Kummer → Frobenius → Schur  
→ Brauer → Fong → Olsson → Külshammer → me → Dauter?

My preferences:

groups

algebras

finite

infinite

# Benjamin Sambale

TU Kaiserslautern

Research Member (GRTA)

Academic genealogy:

... → Gauß → Bessel → Scherk → Kummer → Frobenius → Schur  
→ Brauer → Fong → Olsson → Külshammer → me → Dauter?

My preferences:

- |  |                                   |
|--|-----------------------------------|
| <input checked="" type="checkbox"/> groups | <input type="checkbox"/> algebras |
| <input checked="" type="checkbox"/> finite | <input type="checkbox"/> infinite |
| <input type="checkbox"/> representations   | <input type="checkbox"/> modules  |



# Benjamin Sambale

TU Kaiserslautern

Research Member (GRTA)

Academic genealogy:

... → Gauß → Bessel → Scherk → Kummer → Frobenius → Schur  
→ Brauer → Fong → Olsson → Külshammer → me → Dauter?

My preferences:

- |  |  |
|--|--|
| <input checked="" type="checkbox"/> groups | <input type="checkbox"/> algebras              |
| <input checked="" type="checkbox"/> finite | <input type="checkbox"/> infinite              |
| <input type="checkbox"/> representations   | <input type="checkbox"/> modules               |
|  | <input checked="" type="checkbox"/> characters |

# Benjamin Sambale

TU Kaiserslautern

Research Member (GRTA)

Academic genealogy:

... → Gauß → Bessel → Scherk → Kummer → Frobenius → Schur  
→ Brauer → Fong → Olsson → Külshammer → me → Dauter?

My preferences:

- |  |                                   |
|--|-----------------------------------|
| <input checked="" type="checkbox"/> groups | <input type="checkbox"/> algebras |
| <input checked="" type="checkbox"/> finite | <input type="checkbox"/> infinite |
| <input type="checkbox"/> representations   | <input type="checkbox"/> modules  |
| <input type="checkbox"/> char = 0          | <input type="checkbox"/> char > 0 |
- characters

# Benjamin Sambale

TU Kaiserslautern

Research Member (GRTA)

Academic genealogy:

... → Gauß → Bessel → Scherk → Kummer → Frobenius → Schur  
→ Brauer → Fong → Olsson → Külshammer → me → Dauter?

My preferences:

- |  |  |
|--|--|
| <input checked="" type="checkbox"/> groups   | <input type="checkbox"/> algebras            |
| <input checked="" type="checkbox"/> finite   | <input type="checkbox"/> infinite            |
| <input type="checkbox"/> representations     | <input type="checkbox"/> modules             |
| <input checked="" type="checkbox"/> char = 0 | <input checked="" type="checkbox"/> char > 0 |
- characters

# Benjamin Sambale

TU Kaiserslautern

Research Member (GRTA)

Academic genealogy:

... → Gauß → Bessel → Scherk → Kummer → Frobenius → Schur  
→ Brauer → Fong → Olsson → Külshammer → me → Dauter?

My preferences:

- |  |  |
|--|--|
| <input checked="" type="checkbox"/> groups   | <input type="checkbox"/> algebras              |
| <input checked="" type="checkbox"/> finite   | <input type="checkbox"/> infinite              |
| <input type="checkbox"/> representations     | <input type="checkbox"/> modules               |
| <input checked="" type="checkbox"/> char = 0 | <input checked="" type="checkbox"/> char > 0   |
| splitting fields?                            | <input checked="" type="checkbox"/> characters |

# Benjamin Sambale

TU Kaiserslautern

Research Member (GRTA)

Academic genealogy:

... → Gauß → Bessel → Scherk → Kummer → Frobenius → Schur  
→ Brauer → Fong → Olsson → Külshammer → me → Dauter?

My preferences:

- |  |  |
|--|--|
| <input checked="" type="checkbox"/> groups   | <input type="checkbox"/> algebras            |
| <input checked="" type="checkbox"/> finite   | <input type="checkbox"/> infinite            |
| <input type="checkbox"/> representations     | <input type="checkbox"/> modules             |
| <input checked="" type="checkbox"/> char = 0 | <input checked="" type="checkbox"/> char > 0 |
| splitting fields?                            | Yes, please!                                 |
- characters

## My favorite problem

Let  $G$  be a finite group and  $p$  a prime.

## My favorite problem

Let  $G$  be a finite group and  $p$  a prime.

Let  $B \subseteq \text{Irr}(G)$  be a  $p$ -block and let

$$p^d := \frac{|G|_p}{\min\{\chi(1)_p : \chi \in B\}}.$$

## My favorite problem

Let  $G$  be a finite group and  $p$  a prime.

Let  $B \subseteq \text{Irr}(G)$  be a  $p$ -block and let

$$p^d := \frac{|G|_p}{\min\{\chi(1)_p : \chi \in B\}}.$$

Richard Brauer, 1946:



## My favorite problem

Let  $G$  be a finite group and  $p$  a prime.

Let  $B \subseteq \text{Irr}(G)$  be a  $p$ -block and let

$$p^d := \frac{|G|_p}{\min\{\chi(1)_p : \chi \in B\}}.$$

Richard Brauer, 1946:

**THEOREM 8:** *A block  $B$  of defect  $d$  contains at most  $p^{d(d+1)/2}$  ordinary characters.*

It is probable that the bound  $p^{d(d+1)/2}$  here can be replaced by  $p^d$ , but I have been able to prove this stronger result only for  $d = 0, 1, 2$ .

## My favorite problem

Let  $G$  be a finite group and  $p$  a prime.

Let  $B \subseteq \text{Irr}(G)$  be a  $p$ -block and let

$$p^d := \frac{|G|_p}{\min\{\chi(1)_p : \chi \in B\}}.$$

Richard Brauer, 1946:

**THEOREM 8:** *A block  $B$  of defect  $d$  contains at most  $p^{d(d+1)/2}$  ordinary characters.*

It is probable that the bound  $p^{d(d+1)/2}$  here can be replaced by  $p^d$ , but I have been able to prove this stronger result only for  $d = 0, 1, 2, \mathbf{3}$